

## DISTRIBUTION OF ABO AND RH BLOOD GROUP ALLELES IN GUJRAT REGION OF PUNJAB, PAKISTAN

\*Mohammad Anees and Mohammad Shabir Mirza

*Department of Biological Sciences, The University of Gujrat, G.T. Road College, Gujrat, Pakistan and Aziz Bhatti Shaheed Hospital, Gujrat, Pakistan.*

Received August 2005, accepted November 2005

Communicated by Prof. Dr. M. A. Hafeez

**Abstract:** A study has been carried out on blood groups, representing a random population sample from urban and rural areas of Gujrat region, Province of the Punjab, Pakistan. Blood samples were collected from the patients visiting the Aziz Bhatti Shahid Hospital, Gujrat. The objective of this study was to determine the frequency of different blood groups and their alleles in this region, which would not only help in blood transfusion services, but also eliminate the risk of erythroblastosis and some other diseases. Blood grouping was carried out over a period of 12 months from January 2004 to December 2004, and encompassed 2647 subjects, in which 80.59% were male and 19.41% female. The blood groups were categorized according to ABO/Rh system and allele frequency was determined according to Hardy-Weinberg law. The distribution of phenotypes in the total sample was 0.2489, 0.3691, 0.0688 and 0.3132 for groups A, B, AB and O respectively, with Rh-positive (R) 0.7958 and Rh-negative (r) 0.2042. The distribution of the alleles in the total sample was 0.1740, 0.2229, 0.0435 and 0.5596, for IA, IB, IAIB and i, respectively. From these studies it was concluded that phenotypically B group was dominant in the Gujrat region, with a high allelic frequency of O group.

**Keywords:** Alleles, gene frequency, blood groups, Rh factor, transfusion

### Introduction

The regulation of ABO blood group system is under the control of ABO gene expression [1]. A loss of blood group A antigen expression has been reported in Bladder cancer, caused by the allelic loss or methylation of ABO gene [2]. Research on ABO group system has been of immense interest, due to its medical importance in different diseases. The ABO blood group system is not only important in blood transfusions, cardiovascular diseases, organ transplantation, erythroblastosis in neonates, but also one of the strongest predictors of national suicide rate and a genetic marker of obesity [3-8]. A significant deficit of group O has suggested that there may be susceptibility to develop osteoarthritis in normal hip-joint and spinal osteochondrosis [9,10].

The genetic history of a person can be known by studying the blood groups [11]. For instance, type O blood is the oldest blood and shows a connection to the hunter-gatherer cultures. This blood type is strongly aligned with high animal protein consumption; individuals generally produce higher stomach acids and experience more incidence of gastric ulcer disease than the other groups. Blood group A is primarily associated with vegetarian food sources and individuals in this group secrete smaller amounts of stomach acid and have lesser chances for gastric ulcers, heart diseases, cancer and diabetes [12].

The carbohydrate antigens A, B and O appear to be located on the long arm of the autosomal locus at chromosome number 9 [13], which constitute the four blood types. The gene symbols IA, IB, IAIB and i are often used to denote these alleles. Two alleles, R and r, are responsible for the inheritance

\*Author for correspondence.

E-mail address: anees\_4\_9@hotmail.com



earlier studies. The distribution of the alleles in the total sample was 0.1740, 0.2229, 0.0435 and 0.5596 for IA, IB, IAIB and i respectively, indicating that the allelic frequency of O positive group was higher than the other blood groups. Table 3 compares the distribution of allele frequencies of Rh factor antigens in the Gujrat population with earlier studies on different populations. The distribution of Rh alleles was 0.7958 and 0.2042 for R and r gene, suggesting the dominance of Rh positive. The studied population exhibited the frequency in the order of  $i > IB > IA > IAIB$  and  $R > r$  respectively.

## Discussion

In the study under discussion, the relative

frequency of the various blood group alleles does not seem to deviate from those which have been recorded in the past for most of the Pakistani regions (Table 2). This study has demonstrated that phenotypically, there was a dominance of B blood group ( $B=0.3652$ ,  $O=0.3322$ ,  $A=0.2475$ ), but the allele frequency of O blood group was higher ( $O=0.5596$ ,  $A=0.1901$ ,  $B=0.2503$ ), than the other blood groups. These findings are inconsistent with the results obtained in earlier studies carried out in Wah Cantt ( $O=0.5400$ ,  $A=0.1813$ ,  $B=0.2450$ ) and Punjab division ( $O=0.5760$ ,  $0.2440$ ,  $B=0.1380$ ), but significantly higher from the other regions of Pakistan (Bannu,  $O=0.2507$ ,  $A=0.3103$ ,  $B=0.3623$ , Peshawar  $O=0.3100$ ,  $A=0.2800$ ,  $B=0.3400$ , Swabi,  $O=0.3220$ ,  $A=0.2760$ ,  $B=0.3040$ , Hazara,

**Table 2.**  
Allelic frequency of blood groups (ABO) studied in different populations.

Populations	Frequency of blood groups				References
	A	B	AB	O	
American Indian	0.0390	0.0110	0.000	0.9500	16
Turky	0.1220	0.1213	0.0085	0.7398	17
Nairobi (Kenya)	0.1580	0.1261	0.0239	0.6900	15
Sudan	0.1814	0.1235	0.0268	0.6683	18
Kuwait	0.1608	0.1400	0.0265	0.6678	19
W.Germany	0.2565	0.081	0.0225	0.6400	20
Ukraine	0.2360	0.2250	0.0704	0.5760	21
Hungary	0.2766	0.1218	0.0423	0.5593	22
Nigeria	0.2443	0.2388	0.0275	0.4894	23
Kenya	0.2620	0.2200	0.0440	0.4748	15
Britain (UK)	0.4170	0.0860	0.0300	0.4670	25
India	0.2470	0.3750	0.0530	0.3250	25
Bannu (Pakistan)	0.3103	0.3623	0.0767	0.2507	26
Peshawar (Pakistan)	0.2800	0.3400	0.0700	0.3100	27
Swabi (Pakistan)	0.2760	0.3040	0.0880	0.3220	28
Hazara (Pakistan)	0.2400	0.3200	0.1100	0.3300	29
Bahawalpur (Pakistan)	0.2100	0.3600	0.0600	0.3700	30
Wah cant (Pakistan)	0.1813	0.2450	0.0517	0.5400	31
Punjab (Pakistan)	0.2440	0.1380	0.0420	0.5760	32
<b>Gujrat (Pakistan)</b>	<b>0.1740</b>	<b>0.2229</b>	<b>0.0435</b>	<b>0.5596</b>	<b>Present study</b>

**Table 3.**  
Frequency of Rh antibody allele in different populations.

Populations	Allele frequency		References
	R	r	
Nigeria	0.9430	0.0570	23
Azad Jammu and Kashmir	0.8480	0.1520	33
Kenya	0.8030	0.1970	24
Sudan	0.7436	0.2564	18
Bannu (Pakistan)	0.6720	0.3280	26
Lahore (Pakistan)	0.7170	0.2830	34
Islamabad (Pakistan)	0.7290	0.2710	35
Wah Cantt (Pakistan)	0.7300	0.2710	31
Peshawar (Pakistan)	0.7680	0.2320	27
<b>Gujrat (Pakistan)</b>	<b>0.7958</b>	<b>0.2042</b>	<b>Present study</b>

O=0.3300, A=0.2400, B=0.3200, Bahawalpur, O=0.3700, A=0.2100, B=0.3600). However, the data for the American Indians, Turkey, Nairobi (Kenya), Sudan, Kuwait, W.Germany, Ukraine, Hungary, Nigeria, Kenya and Britain, presented in the same Table, reveals that there is dominance of O group, in these populations in contrast to the Indo-Pak sub-continent, in which both B and O groups show comparable frequency. The least reported group in all the populations has been AB. These studies suggest that the heterogeneity in these populations is due to the genetic and environmental factors which are responsible for varying frequency of the blood groups [26]. Perhaps the environment of the Indo-pak region helps in the expression of blood group B alleles, which are progressively dominating in this region, rather than the oldest O blood group.

In terms of presence of Rh antibody alleles, the data from several studies on Pakistani as well as African, British and other countries populations is compared in Table 3. The present study has shown 0.7958 Rh positive alleles and 0.2042 Rh negative cases, which are very close to those for Kenya (R=0.8030 r=0.1970) and Peshawar (Pakistan,

R=0.7680, r=0.2320) populations, suggesting that there is no significant difference between the frequencies for these populations. However the frequency of Rh positive alleles was less than the Nigerian population (R= 0.9430, r=0.0570). These findings suggest that Gujrat population seems to show the highest rate of Rh positive alleles compared to the other regions of Pakistan studied so far (Table 3).

It is well known that blood groups are associated with several diseases, like cardiovascular, obesity, osteoarthritis, erthroblastosis in neonates and many other diseases especially, osteoporosis and suicide rate in different nations [2-12]. The prevalence of osteoporosis in the proximal femur and lumbar spine averaged 2.3- and 1.7-fold higher in women with blood type AB than in those with blood type O. Thus, ABO blood group status seems to have a significant relationship to the prevalence of osteoporosis in postmenopausal women [36]. [Lester](#) [7] has recently demonstrated that Blood types are one of the strongest predictors of national suicide rates. He studied that in a sample of 51 nations, the suicide rates were negatively associated with the proportion of people with Type O blood, while

homicide rates were positively associated with A and B blood groups. The data generated in the present study may be useful for health planners, while making efforts to face the future health challenges in the region. In short, generation of a simple database of blood groups, not only provides data about the availability of human blood in case of regional calamities, but also serves to enable insight into possibilities of future burden of diseases.

### Acknowledgements

I wish to express my gratitude to Mrs Naseem, Incharge Blood Bank, Aziz Bhatti Shahid Hospital, Gujrat, for providing help in collection of blood group data.

### References

1. **Kominato, Y., Hata, Y., Matsui, K., and Takizawa, H.** 2005. Regulation of ABO gene expression. *Leg. Med.* 3:371-8511.
2. **Chihara, Y., Sugano, K., Kobayashi, A., Kanai, Y., Yamamoto, H., Nakazono, M., Fujimoto, H., Kakizoe, T., Fujimoto, K., Hirohashi, S. and Hirao, Y.** 2005. Loss of blood group A antigen expression in bladder cancer caused by allelic loss and/or methylation of the ABO gene. *Lab Invest.* 9:158-167.
3. **Molison, P.L.** 1979. *Blood transfusion in clinical medicine.* 6th edition, Blackwell Scientific Publication, Oxford, U.K. pp 239-666.
4. **Egawa, H., Oike, F., Buhler, S., Minamiguchi, S., Haga, H., Uryuhara, K., Kiuchi, T., Kaihara, S. and Tanaka, K.** 2004. Impact of recipient age on outcome of ABO-incompatible living-donor liver transplantation. *Transplantation* 15:403-11.
5. **Shamim, A., Hafeez, M.A. and Ahmad, M.M.** 2002. ABO and Rh blood groups I: Markers of cardiovascular risk and association with lipids and other related risk co variables in a Pakistani population. *Proc. Pakistan Acad. Sci.* 39:47-66.
6. **Komar-Szymborska, M., Szymborski, J., Sleboda, A., Bajkacz, M. and Cioch, E.** 1993. RH and ABO incompatibility in newborns treated in a pediatric hospital. *Wiad Lek.* 46:644-50.
7. **Lester, D.** 2005. Predicting suicide in nations. *Suicide Res.* 9:219-23.
8. **Hein, H.O., Suadicani, P. and Gyntelberg, F.** 2005. The Lewis blood group-a new genetic marker of obesity. *Int. J. Obes. Relat. Metab. Disord.* 29:540-52.
9. **Lourie, J.A.** 1983. Is there an association between ABO blood groups and primary osteoarthritis of the hip? *Ann. Hum. Biol.* 10:381-383.
10. **Nevropatol, Z.P.** 1979. Analysis of the distribution of ABO system blood groups among patients with spinal osteochondrosis syndrome. *Ann. Hum. Biol.* 79:409-413.
11. **Sokolov, R.** 1993. *Why We Eat What We Eat: How Columbus Changed the Way the World Eats.* New York: Simon & Schuster. pp 1-50.
12. **Viola, H., J. and Carolyn, M.** 1991. *Seeds of change: five hundred years since Columbus.* 2nd ed., Washington and London, Smithsonian Institution. pp 110-153.
13. **Hoffbrand, A.V.** 1981. *Postgraduate hematology.* 2nd edition, Heinemann Professional Publishing Ltd., London, U.K. pp. 270-350.
14. **Strickberger, M.W.** 1976. *Dominance relations and multiple alleles in diploid organisms.* 2<sup>nd</sup> edition, McMillan Publishing Company Inc., New York, N.Y., U.S.A. pp. 164-180.
15. **Lyko, J., Gaertner, H., Kaviti, J.N., Karithi, M.W. and Akoto, B.** 1992. The blood groups antigens ABO and Rh in Kenyans. *Hamdard Medicus* 35:59-67.
16. **Mourant, A.E., Kipiec, A.C., and Domanjewska, S. K.** 1976. *The distribution of human blood groups.* Oxford University Press, London. pp.10-19.
17. **Akbas, F., Aydin, M. and Cenani, A.** 2003. ABO blood subgroup allele frequencies in the Turkish population. *Anthropol. Anz.* 61:257-60.
18. **Khalil, I.A., Phrykian, S. and Farr, A.D.** 1989. Blood group distribution in Sudan. *Gene Geogr.* 3:7-10.
19. **Al-Bustan, S., El-Zawahri, M., Al-Azmi, D., Al-Bashir, A.A., Miser, A., Geraci, T.K. and Wennberg, R.P.** 2002. Allele frequencies and molecular genotyping of the ABO blood group system in a Kuwaiti population. *Int. J. Hematol.* 75:147-53.
20. **Wagner, F.F., Kasulke, D., Kerowgan, M. and Flegel, W.A.** 1995. Frequencies of the bloods group ABO, Rhesus D category VI. Kell and of clinically relevant high frequency antigens in southwestern Germany. *Infusions her Transfusions med.* 22:285-290.
21. **Mukhin, V.N., Chinakh, D.G., Avdeev, A.V., Kuleba, V.V. and Afanas'ev, M.V.** 2003. Gene frequencies and heterozygosity of the ABO and RH blood group alleles in the populations of two cities of the Donetsk region, Ukraine. *Rus. J. Genetics* 39:530-43.
22. **Tuaszik, T.** 1995. Heterogeneity in the distribution of ABO blood groups in Hungary. *Gene Geogr.* 9:169-176.
23. **Falusi, A.G., Ademowo, O.G., Latunji, C.A., Okeke, A.C., Olatunji, P.O., Onyekwere, T.O, Jimmy, E.O.,**

- Raji, Y., Hedo, C.C., Otukonyong, E.E. and Itata, E.O.** 2000. Distribution of ABO and Rh genes in Nigeria. *Afr. J. Med. Sci.* 29:23-6.
24. **Lyko, J., Gaertner, H., Kaviti, J.N., Karithi, M.W., Akoto, B.** 1992. The blood groups antigens ABO and Rh in Kenyans. *Hamdard Medicus* 35:59-67.
25. **Talib, V.H.** 1991. *Handbook of medical laboratory technology*. 2nd edition. New Delhi, CBS Publishers, pp. 23-47.
26. **Khan, M.S., Subhan, F., Tahir, F., Kazi, B.M., Dil, A.S., Sultan, S., Deepa, F., Khan, F. and Sheikh, M.A.** 2004. Prevalence of blood groups and Rh factor in Bannu region (NWFP) Pakistan. *Pak. J. Med. Res.* 43:8-10.
27. **PMRC.** 1982. Some normal parameters of Pakistani are in the Peshawar area. Peshawar. *Pak. Med. Res. Council*, KMC. 1976-1982.
28. **Khurshid, B., Naz, M., Hassan, M. and Mabood, S.F.** 1992. Frequency of ABO and Rh (D) blood groups in district Swabi N.W.F.P (Pakistan). *J. Sci. Tech. Univ., Peshawar* 16:5-6.
29. **Khalid, M.A., Khan, J.A., Shah, H. and Khan, S.P.** 1984. Frequency of ABO and Rh (D) blood groups in Hazara Division (Abbottabad). *Pak. J. Med. Res.* 23:102-3.
30. **Yousaf, M., Yousaf, N. and Zahid, A.** 1988. Pattern of ABO and Rh (D) blood group distribution in Bahawalpur Division. *Pak. J. Med. Res.* 27:40-51.
31. **Khan, I.M., Khan, G.A. and Akbar, A.** 1982. Gene frequency of ABO and Rh blood group system in Wah Cantonment. Botany Department, Peshawar University, Pakistan.
32. **Shami, S.A. and Rasmuson, M.** 1994. Genetic heterogeneity and gene diversity in the population of Punjab, Pakistan, based on ABO and Rh (D) blood group frequencies. *Hum. Hered.* 44:214-9.
33. **Rashid, M.** 1983. Gene frequency of ABO, blood groups in Azad Kashmir. MSc thesis, Department of Botany, University of Peshawar, Peshawar, Pakistan.
34. **Parveen, N.** 1983. Incidence of ABO and Rh system in Lahore area. MPhil thesis, University of Peshawar, Peshawar, Pakistan.
35. **Subhan, F., Tahir, F., Sultan, S. and Dil, A.S.** 2000. Bad obstetric history: cytogenetic and haematological parameters. *Pak. J. Med. Res.* 39:139-44.
36. **Jong, W.C. and Soo, H.P.** 2004. Associations between ABO blood groups and osteoporosis in postmenopausal women. *Ann. Clin. Lab. Sci.* 34:150-153.