

LEVERRIER-FADDEEV'S ALGORITHM APPLIED TO SPACETIMES OF CLASS ONE

¹J. López-Bonilla, ¹E. Ramírez-García, ²J. Morales and ²G. Ovando

¹Sección de Estudios de Posgrado e Investigación, Escuela Superior de Ingeniería Mecánica y Eléctrica, Instituto Politécnico Nacional, Edif. Z-4, 3er. Piso, Col. Lindavista, CP 07738 México DF, and ²División de Ciencias Básicas e Ingeniería, Universidad Autónoma Metropolitana-Azcapotzalco, Av. San Pablo 180, CP 02200 México DF

Received October 2005, accepted January 2006

Communicated by Prof. Dr. M. Iqbal Choudhary

Abstract: We explain that the Leverrier-Takeno's method to construct the characteristic polynomial of an arbitrary matrix \tilde{A} , plus the Cayley-Hamilton theorem, it is equivalent to the Faddeev's process to obtain \tilde{A}^{-1} . We apply this algorithm to the second fundamental form of a spacetime embedded into flat 5-space.

Keywords: Cayley-Hamilton theorem, Faddeev algorithm, Leverrier-Takeno method, Riemannian 4-spaces of class one.

Introduction

For any matrix $\tilde{A}_{n \times n} = (A^i_j)$ its characteristic equation:

$$\lambda^n + a_1 \lambda^{n-1} + \dots + a_{n-1} \lambda + a_n = 0 \quad (1)$$

can be obtained, through several procedures [1-5], directly from the condition $\det(A^i_j - \lambda \delta^i_j) = 0$. The approach of Leverrier-Takeno [1,6-9] is a simple and interesting technique to construct (1) based in the traces of the powers \tilde{A}^r , $r=1, \dots, n$.

On the other hand, it is well known that an arbitrary matrix \mathbf{A} satisfies its characteristic polynomial:

$$\tilde{A}^n + a_1 \tilde{A}^{n-1} + \dots + a_{n-1} \tilde{A} + a_n I = 0, \quad (2)$$

which is the Cayley-Hamilton identity. If \tilde{A} is non-singular (that is, $\det \tilde{A} \neq 0$), then from (2) we obtain its inverse matrix:

$$\tilde{A}^{-1} = -\frac{1}{a_n} (\tilde{A}^{n-1} + a_1 \tilde{A}^{n-2} + \dots + a_{n-1} I), \quad (3)$$

where $a_n \neq 0$ because $a_n = (-1)^n \det \tilde{A}$.

Faddeev [10-13] proposed an algorithm to determine \tilde{A}^{-1} in terms of \tilde{A} and their traces. Here we exhibit that (3) coincides with the Faddeev's result if we employ the Leverrier-Takeno's formulae for a_j . After, we apply this analysis to second fundamental form \mathbf{b} governing the extrinsic geometry of Riemannian 4-spaces of class one (that is, 4-spaces embedded into (E_5) [14-18]. The corresponding eq. (3) leads to an original expression for its inverse matrix as function of \mathbf{b} and the double dual of Riemann tensor (projection of it onto the Levi-Civita tensor) [18,19]. Normally the Leverrier-Takeno-Faddeev technique is considered useful only in problems of numerical analysis, but the aim of our work is to show the importance of this algorithm in geometrical theories as general relativity, with potential applications to several physical fields.

Leverrier-Takeno and Faddeev methods

If we define the quantities

$$a_0 = 1, \quad s_r = \text{tr } \tilde{A}^r, \quad r = 1, 2, \dots, n \quad (4)$$

then the process of Leverrier-Takeno[1,6-9] implies (1) wherein the a_i are determined with the recurrence expression

$$\begin{aligned} r a_r + s_1 a_{r-1} + s_2 a_{r-2} + \dots + s_{r-1} a_1 + s_r &= 0, \\ r &= 1, 2, \dots, n \end{aligned} \quad (5)$$

Therefore

$$\begin{aligned} a_1 &= -s_1, & 2! a_2 &= (s_1)^2 - s_2, \\ 3! a_3 &= -(s_1)^3 + 3s_1 s_2 - 2s_3, \end{aligned} \quad (6)$$

$$4! a_4 = (s_1)^4 - 6(s_1)^2 s_2 + 8s_1 s_3 + 3(s_2)^2 - 6s_4, \text{ etc.}$$

In particular $\det \tilde{A} = (-1)^n a_n$, that is, the determinant of any square matrix only depends on the traces s_r , which means that \tilde{A} and its transpose have the same determinant.

The Faddeev procedure [10-13] to obtain \tilde{A}^{-1} is a sequence of algebraic computations on the powers \tilde{A}^r and their traces. In fact, his algorithm is given by the instructions

$$\begin{aligned} \tilde{A}_1 &= \tilde{A}, \quad q_1 = \text{tr } \tilde{A}_1, \quad \tilde{B}_1 = \tilde{A}_1 - q_1 \tilde{I}, \\ \tilde{A}_2 &= \tilde{B}_1 \tilde{A}, \quad q_2 = \frac{1}{2} \text{tr } \tilde{A}_2, \quad \tilde{B}_2 = \tilde{A}_2 - q_2 \tilde{I}, \\ \tilde{A}_{n-1} &= \tilde{B}_{n-2} \tilde{A}, \quad q_{n-1} = \frac{1}{n-1} \text{tr } \tilde{A}_{n-1}, \quad \tilde{B}_{n-1} \\ &= \tilde{A}_{n-1} \end{aligned} \quad (7)$$

$$\tilde{A}_n = \tilde{B}_{n-1} \tilde{A}, \quad q_n = \frac{1}{n} \text{tr } \tilde{A}_n$$

Then

$$\tilde{A}^{-1} = \frac{1}{q_n} \tilde{B}_{n-1} \quad (8)$$

For example, if we apply (7) for $n = 4$, then it is easy to see that the corresponding q_r imply (6) with $q_j = -a_j$, and besides (8) reproduces (3). By mathematical induction one can prove that (7) and (8) are equivalent to (3), (4) and (5), showing thus that the Faddeev technique has its origin in the Leverrier-Takeno method plus Cayley-Hamilton theorem.

Spacetimes of class one

A R_4 can be embedded into E5 (that is, the 4-space has class one) if and only if there exists the second fundamental form $b_{ca} = b_{ca}$ satisfying the Gauss-Codazzi equations [14-18,20]

$$R_{acij} = \hat{a}(b_{ai} b_{cj} - b_{aj} b_{ci}), \quad (9)$$

$$b_{ij;c} = b_{ic;j} \quad (10)$$

where $\hat{a} = \pm 1$, R_{acij} is the Riemann tensor and ; r means covariant derivative. It is well-known [21] that whenever $\det(b^i_j)$ is different to zero then (9) implies (10). In other words, when a non-singular matrix \mathbf{b} satisfies the Gauss equation, the Codazzi equation is verified automatically. However, in general the construction of \mathbf{b} for a given spacetime should involve the study of both (9) and (10) together.

Employing (9) it is not difficult to deduce the following result [14,15,22].

$$-24 \det(b^i_j) = K_2^* R^{*ijac} R_{ijac} \quad (11)$$

where K_2 is a Lanczos invariant [19,23-25] defined in terms of the double dual of curvature tensor

$${}^* R^{*ij}_{ac} = \frac{1}{4} \eta^{ijrt} R_{rt}{}^{mn} \eta_{mnac}, \quad (12)$$

with η_{jrt} denoting the Levi-Civita symbol, then the Bianchi identities [20] adopt the compact

form [18,19]

$${}^*R^*{}_{jac;i} = 0, \quad (13)$$

The present work deals with the case $K_2 \neq 0$, thus, according to (11) this implies that the inverse matrix b^{-1}_{ij} exists. In this situation we employ (2), (6), (9) and (11) to obtain the characteristic polynomial of \mathbf{b} [15,16]

$$\mathbf{b}^4 - b\mathbf{b}^3 - \frac{\varepsilon}{2}R\mathbf{b}^2 - p\mathbf{b} - \frac{K_2}{24}\mathbf{I} = \mathbf{0}, \quad (14)$$

where $R = R^i{}_j$ is the scalar curvature and

$$b = tr \mathbf{b}, \quad p = \frac{\varepsilon}{3}b^{ac}G_{ac},$$

$$G_{ij} = {}^*R^*{}_{ija} = \text{Einstein tensor}, \quad (15)$$

Then with (3) and (14) we construct the inverse matrix of \mathbf{b}

$$\frac{K_2}{24} b^{-1}_{ij} = \varepsilon b_{ir} G^r{}_j - p g_{ij}, \quad (16)$$

In a given spacetime we know the metric tensor g_{ij} , G_{ac} and K_2 ; if besides we have \mathbf{b} and ε , then (15) determines p . Thus (16) gives us \mathbf{b}^{-1} and

$$b^{-1r}{}_r = tr \mathbf{b}^{-1} = -\frac{24p}{K_2}. \quad (17)$$

On the other hand, the double dual (12) admits the expansion [16,19,23]

$${}^*R^*{}_{jka} = R_{jka} + G_k g_{ja} + G_a g_{jk} - G_a g_{jc} - G_c g_{ja} + \frac{R}{2}(g_c g_{ja} - g_a g_{jc}), \quad (18)$$

Therefore (9), (16), (17) and (18) imply the original relation

$$K_2 b^{-1}{}_{ia} = \delta a^{\circ} {}^*R^*{}_{ijac} b^{jc}, \quad (19)$$

which means that \mathbf{b}^{-1} essentially is the projection of \mathbf{b} onto double dual tensor. The Codazzi equation (10) is a differential condition on \mathbf{b} .

However, we searched in the literature some differential restriction on \mathbf{b}^{-1} , but without success in this quest. The importance of (19) is that it, (10) and (13) generate such differential requisite

$$(K_2 b^{-1i}{}_a)_{;i} = 0, \quad (20)$$

which is other original contribution of this work.

The deduction of (19) and (20) shows the usefulness of Leverrier-Faddeev-Takeno expressions in the study of 4-spaces of class one. However, our analysis also is applicable to geometric control theory, quantum information processing, field theory, quantum groups and chaos theory (see [26-32]).

References

1. **Wayland, H.** 1945. Expansion of determinantal equations into polynomial form. *Quart. Appl. Math.* 2:277-306.
2. **Householder, A.S. and Bauer, F.L.** 1959. On certain methods for expanding the characteristic polynomial. *Numerische Math.* 1:29-37.
3. **Wilkinson, J.H.** 1965. *The algebraic eigenvalue problem.* Clarendon Press, Oxford.
4. **Lovelock, D. and Rund, H.** 1975. *Tensors, differential forms, and variational principles.* John Wiley and Sons, New York.
5. **Lanczos, C.** 1988. *Applied analysis.* Dover, New York.
6. **Leverrier, U.J.J.** 1840. Sur les variations séculaires des elements elliptiques des sept planets principales. *J. de Math. Pures Appl. Série I* 5:220-254.
7. **Krylov, A.N.** 1931. O cislennom resenii uravnenija, kotorym v techniceskih voprasah opredelajutsja castoty malyh kolebanii material'nyh system. *Izv. Akad. Nauk. SSSR Ser. Fiz.-Mat.* 4:491-539.
8. **Takeno, H.** 1954. A theorem concerning the characteristic equation of the matrix of a tensor of the second order. *Tensor New Series* 3:119-122.
9. **Wilson, E.B., Decius, J.C. and Cross, P.C.** 1980. *Molecular vibrations.* Dover, New York.
10. **Faddeeva, V.N.** 1959. *Computational methods of linear algebra.* Dover, New York.
11. **Faddeev, D.K. and Faddeeva, V.N.** 1963. *Methods in linear algebra.* W. H. Freeman, San Francisco, USA.

12. **Gower, J.C.** 1980. A modified Leverrier-Faddeev algorithm for matrices with multiple eigenvalues. *Linear Algebra Appls.* 31:61-70.
13. **Caltenco, J.H., López-Bonilla, J. and Peña-Rivero, R.** 2000. Characteristic polynomial of A and Faddeev's method for A^{-1} . *Aligarh Bull. Math.* 19:55-59.
14. **López-Bonilla, J. and Núñez-Yépez, H.N.** 1996. An identity for spacetimes embedded into E_5 . *Pramana J. Phys.* 46:219-221.
15. **López-Bonilla, J., Morales, J. and Ovando, G.** 2000. An identity for spacetimes embedded into E_5 . *Indian J. Math.* 42: 09-312.
16. **Caltenco, J.H., López-Bonilla, J. and Ovando, G.** 2001. Spacetime embedded into E_5 . *J. Bangladesh Acad. Sci.* 25:95-97.
17. **Caltenco, J.H., Linaresy, M.R. and López-Bonilla, J.** 2005. A note on the potential for the Gödel solution. *Proc. Pakistan Acad. Sci.* 42:153-154.
18. **Acevedo, M., López-Bonilla, J. and Vidal-Beltrán, S.** 2005. Some applications of a Lovelock's theorem. *Proc. Pakistan Acad. Sci.* 42:249-252.
19. **Lanczos, C.** 1962. The splitting of the Riemann tensor. *Rev. Mod. Phys.* 34:379-389.
20. **Kramer, D., Stephani, H., MacCallum, M. and Herlt, E.** 1980. *Exact solutions of Einstein's field equations.* Cambridge University Press, England.
21. **Thomas, T.Y.** 1936. Riemann spaces of class one and their characterizations. *Acta Math.* 67:169-211.
22. **Fuentes, R., López-Bonilla, J., Matos, T. and Ovando, G.** 1989. Spacetime of class one. *Gen. Rel. Grav.* 21:777-784.
23. **Lanczos, C.** 1938. A remarkable property of the Riemann-Christoffel tensor in four dimensions. *Ann. Math.* 39:842-850.
24. **Gaftoi, V., López-Bonilla, J. and Ovando, G.** 1998. The Lanczos invariants as ordinary divergences. *Nuovo Cim. B* 113:1489-1492.
25. **López-Bonilla, J., Ovando, G. and Puente, O.** 2002. Lanczos scalars as exact divergences. *Indian J. Theor. Phys.* 50:177-180.
26. **Urrutia, L. and Morales, N.** 1994. The Cayley-Hamilton theorem for supersymmetries. *J. Phys. A: Math. Gen.* 27:1981-1997.
27. **Pèoux, G. Monnerville, M. and Flament, J-P.** 1996. Exact analytical polynomial formulation of the exponential of fully matrices. *J. Phys. B: At. Mol. Opt. Phys.* 29: 6031-6047.
28. **Sudbery, A.** 2001. On local invariants of pure three-qubit states. *J. Phys. A: Math. Gen.* 34: 643-652.
29. **Núñez-Yépez, H.N. and Salas-Brito, A.L.** 2001. Jacobi equations using variational principles. *Phys. Lett. A* 275:218-222.
30. **Fellouris, A.G. and Matiadou, L.K.** 2002. On the minimum polynomial of supersymmetries. *J. Phys. A: Math. Gen.* 35:9183-9197.
31. **Delgado, J., Núñez-Yépez, H.N. and Salas-Brito, A.L.** 2004. On the Lagrangian of variational equations of Lagrangian dynamical systems. *Chaos, Solitons and Fractals* 20:925-935.
32. **Cirilo-Lombardo, D.J.** 2005. Non-Abelian Born-Infeld action, geometry and supersymmetry. *Class. Quantum. Grav.* 22:4987-5004.