



# Impact of *Stevia rebaudiana* Culturing in Liquid Medium: Elevation of Yield and Biomass, Mitigation of Steviol Glycosides

Rabia Javed<sup>1,2\*</sup>, and Buhara Yücesan<sup>3</sup>

<sup>1</sup>Department of Biotechnology, Faculty of Biological Sciences,  
Quaid-i-Azam University, Islamabad 45320, Pakistan

<sup>2</sup>Department of Biology, Faculty of Sciences, Bolu Abant Izzet Baysal University,  
Bolu 14030, Turkey

<sup>3</sup>Department of Seed Science & Technology, Faculty of Natural & Agricultural Sciences,  
Bolu Abant Izzet Baysal University, Bolu 14030, Turkey

**Abstract:** An efficient micropropagation system is developed by culturing nodal segments containing the axillary shoot buds on liquid Murashige and Skoog (MS) medium. Direct shoot and root formation are highly effective in both solid and liquid MS media without any plant growth regulators (PGRs). Interestingly, a significant difference in yield is obtained between solid and liquid cultures. It is revealed that a relatively higher amount of plant biomass is obtained after culturing for 4 weeks in a liquid MS medium. However, the shoots produced on solid MS medium produce a remarkable decline in all physiological parameters. On contrary, the bioactive steviol glycosides (rebaudioside-A and stevioside) content is higher in shoots grown in solid MS on a comparative basis, which could be compensated by higher yield. After hardening off, all the regenerants are effectively grown in the field with a negligible loss (<1 %), and steviol glycosides spectra is again obtained by conducting high-performance liquid chromatography (HPLC) analysis after 10 weeks of the plantation. This method has great potential to be applied on large scale in bioreactors.

**Keywords:** *Stevia rebaudiana*, Micropropagation, Liquid Culturing, Steviol Glycosides

## 1. INTRODUCTION

*Stevia rebaudiana*, belonging to the family Asteraceae, has common names viz. sweet leaf or candy leaf [1]. It is a perennial, bushy shrub native to Northeast Paraguay [2]. *Stevia* is of great commercial importance all around the globe because of the presence of steviol glycosides (SGs) in its leaves that play a vital role in reducing high blood pressure, diabetes (Type II), obesity, and issues related to dentistry [3, 4]. This naturally occurring zero-calorie sweetening compound is involved in the regulation of glucose and insulin in the blood which is crucial in maintaining a balanced diet for the well-being of humans [5]. The major steviol glycosides (SGs) responsible for approximately 300 times more sweetness than sucrose include rebaudioside A (Reb-A), stevioside (ST), and rebaudioside C (Reb-C) [6]. SGs are the potential

bioactive compounds that have been approved in various countries because of their tremendous activity against microbes, cancer, inflammation, and diabetes [4, 7].

*S. rebaudiana* is very beneficial to plants both economically and commercially. Conventional methods for *Stevia* propagation involving vegetative propagation through stem cuttings are very slow. Another big hurdle is that *Stevia* seeds are very tiny and poor in germination [8]. The efficacious production of *Stevia* is limited by these factors, therefore, the adoption of modern techniques of biotechnology is the only solution. Tissue culture is one such biotechnological approach having the ability to mass propagate *Stevia* in a minimum period. In this process, true-to-type progeny is produced from a single mother plant [9]. Recently, few abiotic stress elicitors in appropriate

concentrations have been found very useful in this regard [10-18] that alter the biosynthetic pathway for SGs production [19]. Future challenges involve the formation of plenty of biomass by employing a liquid suspension culture technique for explant cultivation. In this respect, the measurement of physiological parameters of plant growth in liquid culture media is immensely important. On the other hand, seed quality may be difficult to obtain as very often, and there is very low germination frequency, sometimes no germination at all.

Based on our literature survey, since the *Stevia* is a photosensitive plant for the SGs accumulation, only limited studies regarding clonal propagation of *S. rebaudiana* with their field performance are reported. Therefore, the current study in *S. rebaudiana* aims to establish a feasible and effective tissue culture protocol for increasing the mass proliferation of nodal explants in a liquid suspension medium. Moreover, the focus of our study is a comparison of different physiological parameters and SGs of commercial significance in *in vitro* and *ex vitro* (field) plants after being harvested.

## 2. MATERIAL AND METHODS

*S. rebaudiana* seeds were obtained from an agricultural company, Polisan Tarim, Istanbul, Turkey. The washing of seeds was carried out using distilled water and 0.1 % (w/v) mercuric chloride ( $\text{HgCl}_2$ ) to get rid of disinfectants. Then the seeds were kept for germination, after which, the leaf nodes were removed to be used as explants to perform *in vitro* experiments. Murashige and Skoog (MS) medium [20] and 3 % (w/v) sucrose was used to prepare the culture medium. The pH of the medium was adjusted to 5.7–5.8 and then 0.8 % (w/v) plant agar was added for solidification. After autoclaving all media, shoot nodes present in culture plates were incubated in a 16 h light: 8 h dark photoperiod at a relative humidity of 55–60 % and  $24 \pm 1$  °C temperature. The stock plants were obtained for further studies.

In the meanwhile, *in vitro* cultivation was also employed using axillary shoot nodes in liquid MS medium without any PGRs. The triplicate experiment was performed and all flasks were kept in a shaking incubator at  $24 \pm 1$  °C in 16 h/8 h photoperiod for 4 weeks. The data regarding the

mean length of roots and shoots, mean number of leaves and nodes per explant, and dry and fresh biomass were recorded. Afterward, shoots were shifted to glass jars (Magenta B-cap, Sigma-Aldrich, USA) having medium and maintained in the growth room.

Plantlets with healthy roots were taken out from jars and then placed into plastic trays ( $12 \times 9$  cells) having soil: compost: vermiculite: perlite mixture (10:10:1:1 (v/v/v/v)). After acclimatization in portable greenhouse ( $50 \times 70 \times 140$  cm) conditions provided with cool-white fluorescent light ( $50 \mu\text{mol m}^{-2}\text{s}^{-1}$  irradiance) for 4 weeks, well-growing plantlets were moved to the field ( $200 \text{ m}^2$ ), in Gebze, Turkey to carry out field trials for 10 weeks. Field trials were conducted with 10 plants per  $\text{m}^2$  without the application of any fertilizer.

SGs analysis was conducted using leaves from *in vitro* regenerated plantlets and shoots. Plant material taken out from plates and jars was cleaned with water carefully and dried in the oven for 48 h at 60 °C. Later on, dried plant material was grounded into a fine powder and each sample (20mg) was taken in 2 mL centrifuge tube. Incubation was done at 55 °C for 15 min in an ultrasonic bath, after which, samples were centrifuged for 10 min at 12,000 rpm and 25 °C in a centrifuge machine. 0.22  $\mu\text{m}$  PTFE Millipore syringe filters were used for filtering the supernatant that was later on transferred to HPLC tubes for analysis.

The standards (rebaudioside A and stevioside) were prepared by dissolving 1 mg of standard glycosides in 30 % of distilled water and 70 % of acetonitrile. Chromatography was performed with an autosampler, a binary pump solvent delivery system, and a dual-wavelength absorbance detector operating at 350 and 210 nm. The column, with  $150 \times 4.6$  mm length and 5  $\mu\text{m}$  particle size was kept warm in a column oven system at 40 °C. Finally, the isocratic flow was performed using acetonitrile and 1 % (w/v) phosphoric acid buffer mixture at a ratio of 68:32 for 20 min.

All micropropagation experiments were performed in triplicate with the randomized design. Statistical analysis of data was done using SPSS software. ANOVA determined the statistical difference and Duncan's multiple range


test calculated the significance of the difference between means  $\pm$  SE value at a confidence interval of 95 %.

### 3. RESULTS AND DISCUSSION

Interestingly, all physiological parameters were significantly higher in the liquid medium than the nodal explant regeneration in the solid medium. Under these circumstances, several parameters including the mean length of shoots (compare  $8.5 \pm 1.1$  and  $5.0 \pm 0.9$  cm in green bars of Figure 1), mean length of roots (compare  $1. \pm 0.5$  and  $1.1 \pm 0.2$  cm in blue bars of Fig.1) and mean number of roots ( $4.2 \pm 0.8$ ), nodes ( $7.44 \pm 1.5$ ) and leaves ( $17.5 \pm 2.6$ ) were found to be higher than those produced from a solid medium with a mean of  $1.6 \pm 0.1$  roots,  $14.0 \pm 0.8$  leaves, and a mean of  $6.0 \pm 0.4$  nodes as shown in Figure 2. In this study, it was clearly shown that shoot organogenesis in a liquid medium can prove to be very cost-efficient to produce a large number of plants and to obtain excellent yield in a bioreactor system

once established. Similarly, some earlier reports on temporary immersion [21] and bioreactor systems [22] claim the efficiency of liquid culture on a large number of plantlet formations. However, tissue browning during cultivation might be a drawback due to exogenously added growth regulators and that should be overcome by a separate multiplication medium [21]. Long exposure (more than 6 weeks) to the plant tissues in liquid culture might be incorporated with morphological and physiological abnormalities due to the presence of growth regulators [22]. Although some reports resolve using less constant immersion frequencies, well-rooted shoots increased fidelity in field conditions. However, many of these efforts are not available for large-scale production because of the additional passages of callus tissue needed for clonal propagation from microshoots. In the present study, we designed an effective regeneration system with a short regeneration process without callus formation.

Fresh biomass (0.67 g) and dry biomass (0.10 g)



**Fig. 1.** Comparison of solid and liquid media in terms of shoot and root length for 4 weeks of cultivation. Bar lines represent the standard error of the mean values.

were also higher in MS liquid culturing system after 4 weeks of culturing. In this study, it was clearly shown that shoot organogenesis in a liquid medium can prove to be very cost-efficient to produce a large number of plants and to obtain excellent yield in a bioreactor system once established. Moreover, SGs content was also checked in in vitro plants. It was shown that Reb-A content (3.3 %) was significantly higher in solid MS than in liquid MS (1.4 %) medium. Similarly, ST content was almost doubled in solid MS, i.e., ST content was 0.4 % in leaves of liquid MS, much lower than 0.81 % obtained from the solid MS basal medium depicted in Figure 3. Earlier reports aiming at SGs production through the various cell and shoot cultures with or without elicitor [21, 23, 24] did not provide a reasonable amount for sustainable sweetener compound production unless in vitro raised samples were transferred progressively to ex vitro conditions [25]. Since *S. rebaudiana* is a photoperiod-sensitive plant, SGs accumulation is directly affected by a long day period (>14 h).

Healthy and vigorous shoots were shifted to the glass jars having MS fortified with or without rooting phytohormones, i.e., 0.25 mg/L IAA [25], and allowed to grow for 4 weeks. The percentage of rooting frequency obtained was approximately

89 %. All regenerants were healthy, therefore, shifted to about 500 jars of 450 ml volume filled with 0.4 dm<sup>3</sup> of soil mixture. The jars covered with the thin transparent film were placed in a growth chamber having 16 h/8 h photoperiod for 4 weeks. After hardening off in the greenhouse, almost all plants (> 98 %) were transferred to the field. Our findings revealed that all the regenerants produced in vitro were showing enough vigor to sustain the environmental conditions of the field and the average shoot length was measured to be ~ 62 cm after 10 weeks of the plantation. Figure 4 illustrates each step of the process in detail.

Table 1 shows that the Reb-A content increased from 2.6 % (w/v) in shoots to 3.7 % (w/w) in rooted-shoots, depending on the growth and development. The highest SGs content was observed after harvesting the leaf materials from the field 7.2 % (w/w). A significant difference was observed regarding ST content of 4 weeks old shoots (0.8 %, w/w) and 8 weeks old regenerants (2.0 %, w/w). Furthermore, in contrary to the rise of Reb-A content, ST content did not change and remained the same in field samples. The ratio of Reb-A and ST was obtained highest (3.6) in field-grown plants followed by 4-weeks old shoots (3.3). The ratio was lowest (1.9) in 4-weeks old






Fig. 2. Comparison of the mean number of roots, leaves, and nodes produced in solid and liquid medium for 4 weeks.





**Fig. 3.** Comparison of rebaudioside A (Reb-A) and stevioside (ST) content of the leaves produced in solid and liquid MS medium. Note: Bar lines represent the standard error of the mean values.



**Fig. 4.** Steps of tissue culture experiment conducted in *S. rebaudiana* from in vitro to ex vitro stages: Formation of shoots using nodes (explants) on solid MS medium for 4 weeks (A), Formation of shoots using nodes (explants) on liquid MS medium for 4 weeks (B), shoot and roots formation from liquid culture (C), root formation on MS medium having 0.25 mg/L IBA or absence of PGR (D), adventitious root formation from the shoots (E), hardening off the regenerants in jars filled with soil under plant growth room conditions (F), the general appearance of the potted regenerants one week before field plantation (G), flowering stage of the regenerants in field conditions after 12 weeks of plantation to the field (H).

**Table 1.** Comparison of SGs (rebaudioside A and stevioside) in the leaves of 4 weeks-old shoots, regenerants (rooted-shoots), and field samples ( $\pm$ : standard deviation).

Steviol glycosides	Plant Samples (% DW)*		
	Shoots (4-wk old)	Rooted-shoots (4-wk old)	Field samples (10-wk old)
Rebaudioside A	2.6 $\pm$ 0.2 <sup>c</sup>	3.7 $\pm$ 0.3 <sup>b</sup>	7.2 $\pm$ 0.3 <sup>a</sup>
Stevioside	0.8 $\pm$ 0.0 <sup>b</sup>	2.0 $\pm$ 0.1 <sup>a</sup>	2.0 $\pm$ 0.1 <sup>a</sup>
RebA/ST	3.3	1.9	3.6

\* Same letter in the rows represents no significant difference at a 95 % confidence interval. Data were obtained from dried leaf materials.

regenerants due to higher ST content in them. The most probable reason might be the involvement of critical enzymatic reactions in the biochemical pathway of stevioside synthesis during root formation [26].

#### 4. CONCLUSION

In conclusion, based on this protocol, about 32,500 healthy plantlets of ~16 cm in length can be produced within 7 months to conduct field trials. Most importantly, a significantly higher yield of *Stevia* plantlets has been obtained in MS liquid culture at a small laboratory scale. Commercial production of more SGs from these plantlets still needs further refinement of this technique by developing a suitable bioreactor system. Once an automation system is established for *Stevia* plants containing value-added SGs, it is noteworthy to mention a cost-efficient production targeting both agricultural practices and refinery systems for the natural sweeteners.

#### 5. ACKNOWLEDGEMENTS

The authors are grateful to TUBITAK No. 2216 for providing funds to RJ and TUBITAK No. 114-O 006 for financial support to BY. Research support by the Department of Biology, Bolu Abant İzzet Baysal University, Turkey is also appreciated.

#### 6. CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### 7. DECLARATION

The results of this study are original, the same material is neither published nor under consideration elsewhere,

approval of all authors has been obtained, and in case the article is accepted for publication, its copyright will be assigned to the Pakistan Academy of Sciences

#### 8. REFERENCES

1. A.D. Kinghorn. *Stevia: the genus Stevia*. Taylor and Francis Inc., NY, p. 1–17 (2002).
2. D. Soejarto. Botany of *Stevia* and *Stevia rebaudiana*. Taylor and Francis, In: A. Kinghorn (Ed.), *Stevia: The genus Stevia*. London, New York, p. 18–39 (2002).
3. J.C. Liu, P.K. Kao, P. Chan, Y.H. Hsu, C.C. Hou, G.S. Lien, M.H. Hsieh, Y.J. Chen, and Y.T. Cheng. Mechanism of the antihypertensive effect of stevioside in anesthetized dogs. *Pharmacology* 67: 14 (2003).
4. A. Dey, S. Kundu, A. Bandyopadhyay, and A. Bhattacharjee. Efficient micropropagation and chlorocholine chloride induced stevioside production of *Stevia rebaudiana* Bertoni. *CR Biologies* 336: 17–28 (2013).
5. A.D. Anton, C.K. Martin, H. Han, S. Coulon, W.T. Cefalu, P. Geiselman, and Williamson DA. Effects of *Stevia*, aspartame, and sucrose on food intake, satiety, and postprandial glucose and insulin levels. *Appetite* 55: 37–43 (2010).
6. U. Wölwer-Rieck. The leaves of *Stevia rebaudiana* (Bertoni), their constituents and the analyses thereof: a review. *Journal of Agricultural and Food Chemistry* 60: 886–895 (2012).
7. M. Libik-Konieczny, E. Capecka, M. Tuleja, and R. Konieczny. Synthesis and production of steviol glycosides: recent research trends and perspectives. *Applied Microbiology and Biotechnology* 105: 3883–3900 (2021).
8. A.K. Yadav, S. Singh, D. Dhyani, and P.S. Ahuja. A review on the improvement of *Stevia* [*Stevia*

- rebaudiana* (Bertoni)]. *Canadian Journal of Plant Science* 91: 1–27 (2011).
9. Y. Tamura, S. Nakamura, H. Fukui, and M. Tabata. Clonal propagation of *Stevia rebaudiana* Bertoni by stem-tip culture. *Plant Cell Reports* 3: 183–185 (1984).
  10. R. Javed, M. Usman, B. Yucesan., M. Zia, E. Gurel. Effect of zinc oxide (ZnO) nanoparticles on physiology and steviol glycosides production in micropropagated shoots of *Stevia rebaudiana* Bertoni. *Plant Physiology and Biochemistry* 110: 94–99 (2017a).
  11. R. Javed, A. Mohamed, B. Yucesan, G. Ekrem, R. Kausar, and M. Zia. CuO nanoparticles significantly influence in vitro culture, steviol glycosides, and antioxidant activities of *Stevia rebaudiana* Bertoni. *Plant Cell Tissue Organ Culture* 131: 611–620 (2017b).
  12. R. Javed, B. Yucesan, M. Zia, and E. Gurel. Differential effects of plant growth regulators on physiology, steviol glycosides content, and antioxidant capacity in micropropagated tissues of *Stevia rebaudiana*. *Biologia* 72: 1156–1165 (2017c).
  13. R. Javed, B. Yucesan, and E. Gurel. Hydrogen peroxide-induced steviol glycosides accumulation and enhancement of antioxidant activities in leaf tissues of *Stevia rebaudiana* Bertoni. *Sugar Tech* 20: 100–104 (2018a).
  14. R. Javed, B. Yucesan., M. Zia, and E. Gurel. Elicitation of secondary metabolites in callus cultures of *Stevia rebaudiana* Bertoni grown under ZnO and CuO nanoparticles stress. *Sugar Tech* 20: 194–201 (2018b).
  15. R. Javed, and E. Gurel. Salt stress by NaCl alters the physiology and biochemistry of tissue culture-grown *Stevia rebaudiana* Bertoni. *Turkish Journal of Agriculture and Forestry* 43: 11–20 (2019).
  16. M.A. Ahmad, R. Javed, M. Adeel, M. Rizwan, Q. Ao, and Y. Yang. Engineered ZnO and CuO nanoparticles ameliorate morphological and biochemical response in tissue culture regenerants of Candy leaf (*Stevia rebaudiana*). *Molecules* 25: 1356 (2020a).
  17. M.A. Ahmad, R. Javed, M. Adeel, M. Rizwan, and Y. Yang. PEG 6000-stimulated drought stress improves the attributes of *in vitro* growth, steviol glycosides production, and antioxidant activities in *Stevia rebaudiana* Bertoni. *Plants* 9: 1552 (2020b).
  18. M.A. Ahmad, X. Deng, M. Adeel, M. Rizwan, N. Shakoore, Y. Yang, and R. Javed. Influence of calcium and magnesium elimination on plant biomass and secondary metabolites of *Stevia rebaudiana* Bertoni. *Biotechnology and Applied Biochemistry* (2021).
  19. H. Tavakoli, N. Tavakoli, and F. Moradi. The effect of the elicitors on the steviol glycosides biosynthesis pathway in *Stevia rebaudiana*. *Functional Plant Biology* 46: 787–795 (2019).
  20. T. Murashige, and F.A. Skoog. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiology* 15: 473–497 (1962).
  21. R.V. Sreedhar, L. Venkatachalam, R. Thimmaraju, N. Bhagyalakshmi, M.S. Narayan, and G.A. Ravishankar. Direct organogenesis from leaf explants of *Stevia rebaudiana* and cultivation in bioreactor. *Biologia Plantarum* 52: 355–360 (2008).
  22. K. Jiménez-Quesada. Evaluación del desarrollo *in vitro* de *Stevia rebaudiana* Bertoni con el empleo de sistemas de micropropagación semi-sólido, inmersión temporal RITA® y biorreactor de burbujeo. BSc in biology Thesis School of Biology, Costa Rica Institute of Technology, Costa Rica (in Spanish) (2011).
  23. J. Striedner, F. Czygan, and Braunegg G. Contributions to the biotechnological production of sweeteners from *Stevia rebaudiana* Bertoni: a method for the serial analysis of diterpene glycosides by HPLC. *Acta Biotechnologica* 11: 495–499 (1991).
  24. N. Bondarev, O. Reshetnyak, and A. Nosov. Effects of nutrient medium composition on development of *Stevia rebaudiana* shoots cultivated in the roller bioreactor and their production of steviol glycosides. *Plant Science* 165: 845–850 (2003).
  25. B. Yucesan, A. Mohammed, R. Büyükgöçmen, A. Cevher, O. Kavas, S. Gürel, and E. Gürel. In-vitro and ex-vitro propagation of *Stevia rebaudiana* Bertoni with high Rebaudioside-A content—A commercial scale application. *Scientia Horticulturae* 203: 20–28 (2016).
  26. S.M. Swanson, G.B. Mahady, and C.W.W. Beecher. Stevioside biosynthesis by callus, root, shoot and rooted-shoot cultures. *Plant Cell Tissue Organ Culture* 28: 151–157 (1992).

